

Single spin asymmetries in $\ell p^\uparrow \rightarrow hX$ and $e^-e^+ \rightarrow q\bar{q} \rightarrow h^\uparrow X$

M. Anselmino

*Dipartimento di Fisica Teorica, Università di Torino and
INFN, Sezione di Torino, Via P. Giuria 1, 10125 Torino, Italy*

Abstract

Spin observables may reveal much deeper properties of non perturbative hadronic physics than unpolarized quantities. We discuss here possible origins of single spin asymmetries in DIS, absent in the elementary lepton-quark interactions, and suggest strategies to isolate and understand some non perturbative spin dependence of distribution and fragmentation functions. We also discuss the polarization of hadrons produced in e^+e^- annihilation at LEP and show how final state $q\bar{q}$ interactions may give origin to non zero values of the off-diagonal element $\rho_{1,-1}$ of the helicity density matrix of vector mesons: some predictions are given for K^{*0} , ϕ and D^* in agreement with recent OPAL data. Possible analogous effects in DIS and other processes are suggested.

1 Introduction

The spin properties of hadrons inclusively produced in high energy interactions are related to the fundamental properties of quarks and gluons and to their elementary interactions in a much more subtle way than unpolarized quantities. They test unusual basic dynamical properties and reveal how the usual models for quark distribution and hadronization – successful in predicting unpolarized cross-sections – may not be adequate to describe spin effects. We consider here such two cases, single spin asymmetries in Deep Inelastic Scattering and the polarization of mesons produced in the fragmentation of polarized quarks at LEP.

2 Single spin asymmetries in DIS

We discuss first single spin asymmetries in DIS processes. We start by reminding that single spin asymmetries in large p_T inclusive hadronic reactions are forbidden

in leading-twist perturbative QCD, reflecting the fact that single spin asymmetries are zero at the partonic level and that collinear parton configurations inside hadrons do not allow single spin dependences. However, experiments tell us in several cases, [1, 2] that single spin asymmetries are large and indeed non negligible.

The usual arguments to explain this apparent disagreement between pQCD and experiment invoke the moderate p_T values of the data – a few GeV, not quite yet in the true perturbative regime – and the importance of higher-twist effects. Several phenomenological models have recently attempted to explain the large single spin asymmetries observed in $p^\uparrow p \rightarrow \pi X$ as twist-3 effects which might be due to intrinsic partonic $\mathbf{k}_\perp$ in the fragmentation [3] and/or distribution functions [4]-[6].

Let us consider a process in which one has convincing evidence that partons and perturbative QCD work well and successfully describe the unpolarized and leading-twist spin data, namely Deep Inelastic Scattering (DIS). In particular we shall discuss single spin asymmetries in the inclusive, $\ell N^\uparrow \rightarrow \ell + \text{jets}$ and $\ell N^\uparrow \rightarrow hX$, reactions looking at possible origins of such asymmetries and devising strategies to isolate and discriminate among them [7].

According to the QCD hard scattering picture and the factorization theorem [3, 8, 9] the cross-section for the $\ell N^\uparrow \rightarrow hX$ reaction is given by

$$\frac{E_h d^3 \sigma^{\ell+N, S \rightarrow h+X}}{d^3 \mathbf{p}_h} = \sum_{q; \lambda_{q'}, \lambda'_{q'}, \lambda_h} \int \frac{dx d^2 \mathbf{k}_\perp d^2 \mathbf{k}'_\perp}{\pi z} \quad (1)$$

$$\tilde{f}_{q/N}^{N, S}(x, \mathbf{k}_\perp) \frac{d\hat{\sigma}^{q, P_q}}{d\hat{t}}(x, \mathbf{k}_\perp, \mathbf{k}'_\perp) \rho_{\lambda_{q'}, \lambda'_{q'}}^{q'}(x, \mathbf{k}_\perp, \mathbf{k}'_\perp) \widetilde{D}_{\lambda_h, \lambda_h}^{\lambda_{q'}, \lambda'_{q'}}(z, \mathbf{k}'_\perp).$$

Let us briefly discuss the different quantities appearing in the above equation; more details can be found in Ref. [7]. $\tilde{f}_{q/N}^{N, S}(x, \mathbf{k}_\perp)$ is the quark distribution function, that is the total number density of quarks q with momentum fraction x and intrinsic transverse momentum $\mathbf{k}_\perp$ inside a polarized nucleon N with spin four-vector S .

$d\hat{\sigma}^{q, P_q}/d\hat{t}$ is the cross-section for the $\ell q^\uparrow \rightarrow \ell q$ process, with an unpolarized lepton and an initial quark with polarization P_q , while the final quark and lepton polarization are summed over. Notice that for helicity conserving elementary interactions $d\hat{\sigma}^{q, P_q}/d\hat{t}$ equals the unpolarized cross-section $d\hat{\sigma}^q/d\hat{t}$. $\rho^{q'}$ is the helicity density matrix of the final quark produced in the $\ell q^\uparrow$ interaction and

$$\widetilde{D}_{q, s_q}^h(z, \mathbf{k}_\perp) = \sum_{\lambda_q, \lambda'_q} \rho_{\lambda_q, \lambda'_q}^q \widetilde{D}_{\lambda_h, \lambda_h}^{\lambda_q, \lambda'_q}(z, \mathbf{k}_\perp) \quad (2)$$

describes the fragmentation process of a polarized quark q with spin s_q into a hadron h with helicity λ_h , momentum fraction z and intrinsic transverse momentum $\mathbf{k}_\perp$ with respect to the jet axis. It is simply the inclusive cross-section for the $q^\uparrow \rightarrow hX$

fragmentation process. As it will be shown in the second part we can safely neglect the coherent interactions of the fragmenting quark, as we are not looking at the spin state of the final hadron. The usual unpolarized fragmentation function is given by

$$D_q^h(z) = \frac{1}{2} \sum_{\lambda_q, \lambda_h} \int d^2 \mathbf{k}_\perp \widetilde{D}_{\lambda_h, \lambda_q}^{\lambda_q, \lambda_q}(z, \mathbf{k}_\perp). \quad (3)$$

Similar formulae hold also when the elementary interaction is $\ell q \rightarrow \ell q g$ rather than $\ell q \rightarrow \ell q$: in the latter case two jets are observed in the final state – the target jet and the current quark jet – and in the former case three – the target jet and $q + g$ current jets.

In Eq. (1) we have taken into account intrinsic transverse momenta both in the distribution and the fragmentation process; the $\mathbf{k}_\perp$ dependences are expected to have negligible effects on unpolarized variables for which they are indeed usually neglected, but they can be of crucial importance for some single spin observables, as discussed in Refs. [3]-[6].

We discuss now possible sources of single spin effects in Eq. (1).

$\mathbf{k}_\perp$ effects in fragmentation process [3]

The fragmentation process of a transversely polarized quark into a hadron h (whose polarization is not observed) with fixed z and $\mathbf{k}_\perp$ may depend on the quark spin orientation, provided the quark spin s_q has a component perpendicular to the hadron-quark plane (otherwise any spin dependence would be forbidden by parity conservation). That is, there might be a non zero *quark analysing power* [3]:

$$A_q^h \equiv \frac{\widetilde{D}_{q, s_q}^h(z, \mathbf{k}_\perp) - \widetilde{D}_{q, -s_q}^h(z, \mathbf{k}_\perp)}{\widetilde{D}_{q, s_q}^h(z, \mathbf{k}_\perp) + \widetilde{D}_{q, -s_q}^h(z, \mathbf{k}_\perp)}. \quad (4)$$

By rotational invariance $\widetilde{D}_{q, -s_q}^h(z, \mathbf{k}_\perp) = \widetilde{D}_{q, s_q}^h(z, -\mathbf{k}_\perp)$, which shows immediately how the quark analysing power vanishes for $\mathbf{k}_\perp = 0$.

$\mathbf{k}_\perp$ effects in distribution functions [4]-[6]

A similar idea had been previously proposed in Refs. [4, 5] and later rediscovered in Ref. [6], concerning the distribution functions: that is, the number of quarks (whose spin is not observed) with fixed x and $\mathbf{k}_\perp$ inside a transversely polarized nucleon may depend on the nucleon spin orientation and the function

$$\Delta \tilde{f}_{q/N}^{N^\uparrow}(x, \mathbf{k}_\perp) \equiv \tilde{f}_{q/N}^{N^\uparrow}(x, \mathbf{k}_\perp) - \tilde{f}_{q/N}^{N^\downarrow}(x, \mathbf{k}_\perp) = \tilde{f}_{q/N}^{N^\uparrow}(x, \mathbf{k}_\perp) - \tilde{f}_{q/N}^{N^\uparrow}(x, -\mathbf{k}_\perp) \quad (5)$$

may be different from zero. $\uparrow$ and $\downarrow$ refer to the nucleon spin up or down with respect to the quark-nucleon plane.

In terms of the usual light-cone operator definition of structure functions one has [3, 10]

$$\Delta \tilde{f}_{q/N}^{N\uparrow} = 2 \text{Im} \int \frac{dy^- d\mathbf{y}_\perp}{(2\pi)^3} e^{-ixp^+ y^- + i\mathbf{k}_\perp \cdot \mathbf{y}_\perp} \langle p, - | \bar{\psi}_a(0, y^-, y_\perp) \frac{\gamma^+}{2} \psi_a(0) | p, + \rangle. \quad (6)$$

In Ref. [3] it is argued that such off-diagonal (in the helicity basis) matrix elements are zero due to the time-reversal invariance of QCD, and indeed this is proven by exploiting the time-reversal and parity transformation properties of free Dirac spinors. However, in Ref. [10] it has been shown that this need not be so in chiral models with quark moving in a background of chiral fields.

Both $\widetilde{D}_{q\uparrow}^h(z, \mathbf{k}_\perp) - \widetilde{D}_{q\downarrow}^h(z, \mathbf{k}_\perp)$ and $\tilde{f}_{q/N}^{N\uparrow}(x, \mathbf{k}_\perp) - \tilde{f}_{q/N}^{N\downarrow}(x, \mathbf{k}_\perp)$ can be considered as new fundamental spin and $\mathbf{k}_\perp$ dependent non perturbative functions describing respectively quark fragmentation and distribution properties. In the sequel we shall devise strategies to test their relevance.

Single spin effects in the elementary interactions

As we already mentioned both perturbative QED and QCD at high energy do not allow single helicity flips in the ℓq interactions, so that there cannot be any dependence on the quark polarization in $d\hat{\sigma}^{q,P_q}/d\hat{t}$. Similarly, the perturbative QCD evolution of the distribution and fragmentation functions is not expected to introduce any single spin dependence. We must conclude that the hard elementary interactions are unlikely to introduce any single spin effect: however, this basic QED and QCD property should also be tested.

Let us now describe a set of possible measurements which could single out some of the above mechanisms and test them.

a) $\ell N^\uparrow \rightarrow \ell + 2 \text{ jets}$

Here one avoids any fragmentation effect by looking at the fully inclusive cross-section for the process $\ell N^\uparrow \rightarrow \ell + 2 \text{ jets}$, the 2 jets being the target and current ones; this is the usual DIS, and Eq. (1) becomes

$$\frac{d^2 \sigma^{\ell+N, S \rightarrow \ell+X}}{dx dQ^2} = \sum_q \int d^2 \mathbf{k}_\perp \tilde{f}_{q/N}^{N,S}(x, \mathbf{k}_\perp) \frac{d\hat{\sigma}^{q,P_q}}{d\hat{t}}(x, \mathbf{k}_\perp). \quad (7)$$

In this case the elementary interaction is a pure QED, helicity conserving one, $\ell q \rightarrow \ell q$, and $d\hat{\sigma}^{q,P_q}/d\hat{t}$ cannot depend on the quark polarization. Some spin de-

pendence might only remain in the distribution function, due to intrinsic $\mathbf{k}_\perp$ effects [4]-[6], [10] and we have

$$\begin{aligned} \frac{d^2 \sigma^{\ell N^\uparrow \rightarrow \ell + X}}{dx dQ^2} - \frac{d^2 \sigma^{\ell N^\downarrow \rightarrow \ell + X}}{dx dQ^2} &= \sum_q \int d^2 \mathbf{k}_\perp \\ &\times \Delta \tilde{f}_{q/N}^{N^\uparrow}(x, \mathbf{k}_\perp) \frac{d\hat{\sigma}^q}{d\hat{t}}(x, \mathbf{k}_\perp). \end{aligned} \quad (8)$$

Despite the fact that $\Delta \tilde{f}_{q/N}^{N^\uparrow}$ is an odd function of $\mathbf{k}_\perp$ a non zero value of the above difference – of $\mathcal{O}(k_\perp/\sqrt{Q^2})$, twist 3 – might remain even after integration on $d^2 \mathbf{k}_\perp$ because of the $\mathbf{k}_\perp$ dependence of $d\hat{\sigma}^q/d\hat{t}$, similarly to what happens in $pp^\uparrow \rightarrow \pi X$ [6]. The observation of a non vanishing value of the single spin effect of Eq. (8) would be a decisive test in favour of the mechanism suggested in Refs. [4]-[6] and would allow an estimate of the new function (5).

b) $\ell N^\uparrow \rightarrow h + X$ (2 jets, $\mathbf{k}_\perp \neq 0$)

One looks for a hadron h , with transverse momentum $\mathbf{k}_\perp$, inside the quark current jet; the final lepton may or may not be observed. The elementary subprocess is $\ell q \rightarrow \ell q$ and Eq. (1) yields

$$\begin{aligned} &\frac{E_h d^5 \sigma^{\ell + N^\uparrow \rightarrow h + X}}{d^3 \mathbf{p}_h d^2 \mathbf{k}_\perp} - \frac{E_h d^5 \sigma^{\ell + N^\downarrow \rightarrow h + X}}{d^3 \mathbf{p}_h d^2 \mathbf{k}_\perp} \\ &= \sum_q \int \frac{dx}{\pi z} \Delta_T q(x) \Delta_N \hat{\sigma}^q(x, \mathbf{k}_\perp) \left[\tilde{D}_{q^\uparrow}^h(z, \mathbf{k}_\perp) - \tilde{D}_{q^\uparrow}^h(z, -\mathbf{k}_\perp) \right] \end{aligned} \quad (9)$$

where $\Delta_T q$ is the polarized number density for transversely spinning quarks q and $\Delta_N \hat{\sigma}^q$ is the elementary cross-section double spin asymmetry

$$\Delta_N \hat{\sigma}^q = \frac{d\hat{\sigma}^{\ell q^\uparrow \rightarrow \ell q^\uparrow}}{d\hat{t}} - \frac{d\hat{\sigma}^{\ell q^\uparrow \rightarrow \ell q^\downarrow}}{d\hat{t}}. \quad (10)$$

In Eq. (9) we have neglected the $\mathbf{k}_\perp$ effect in the distribution function, which can be done once the asymmetry discussed in *a*) turns out to be negligible. We are then testing directly the mechanism suggested in Ref. [3] and a non zero value of the l.h.s. of Eq. (9) would be a decisive test in its favour and would allow an estimate of the new function appearing in the numerator of Eq. (4). Notice again that even upon integration over $d^2 \mathbf{k}_\perp$ the spin asymmetry of Eq. (9) might survive, at higher twist order $k_\perp/p_T$, due to some $\mathbf{k}_\perp$ dependence in $\Delta_N \hat{\sigma}^q$.

c) $\ell N^\uparrow \rightarrow h + X$ (2 jets, $\mathbf{k}_\perp = 0$)

By selecting events with the final hadron collinear to the jet axis ($\mathbf{k}_\perp = 0$) one forbids any single spin effect in the fragmentation process. As in the fully inclusive case a) the observation of a single spin asymmetry in such a case would imply single, $\mathbf{k}_\perp$ dependent, spin effects in the distribution functions.

d) $\ell N^\uparrow \rightarrow h + X$ (3 jets, $\mathbf{k}_\perp \neq 0$)

The elementary process is now $\ell q \rightarrow \ell q q$ and one looks at hadrons with $\mathbf{k}_\perp \neq 0$ inside the q current jet. Single spin asymmetries can originate either from single spin effects in the fragmentation process or distribution functions. One should not, in principle, forget possible spin effects in the elementary QCD interaction.

e) $\ell N^\uparrow \rightarrow \ell + 3 \text{ jets}$ or $\ell N^\uparrow \rightarrow h + X$ (3 jets, $\mathbf{k}_\perp = 0$)

These cases are analogous to a) and c) respectively: the measurement eliminates spin effects arising from the fragmentation functions. The only possible origin of a single spin asymmetry would reside in the distribution function. However, if no effect is observed in cases a) and c), but some effect is observed here, then one has to conclude that there must be some single spin effect in the elementary QCD interaction. Utterly unexpectedly, this would question the validity of quark helicity conservation, a fundamental property of pQCD which has never been directly tested.

In summary, a study of single transverse spin asymmetries in DIS could provide a series of profound tests of our understanding of large p_T QCD-controlled reactions.

3 $\rho_{1,-1}(V)$ in the process $e^- e^+ \rightarrow q \bar{q} \rightarrow V + X$

We consider now the spin properties of hadrons produced at LEP. It was pointed out in Refs. [11] and [12] that final state interactions between the q and $\bar{q}$ created in $e^+ e^-$ annihilations – usually neglected, but indeed necessary – might give origin to non zero spin observables which would otherwise be forced to vanish: the off-diagonal element $\rho_{1,-1}$ of the helicity density matrix of vector mesons may be sizeably different from zero [11] due to a coherent fragmentation process which takes into account $q\bar{q}$ interactions. The incoherent fragmentation of a single independent quark leads instead to zero values for such off-diagonal elements.

We present predictions [13] for $\rho_{1,-1}$ of several vector mesons V provided they are produced in two jet events, carry a large momentum or energy fraction $z = 2E_V/\sqrt{s}$, and have a small transverse momentum p_T inside the jet. Our estimates are in agreement with the existing data and are crucially related both to the presence of final state interactions and to the Standard Model couplings of the elementary

$e^-e^+ \rightarrow q\bar{q}$ interaction.

The helicity density matrix of a hadron h inclusively produced in the two jet event $e^-e^+ \rightarrow q\bar{q} \rightarrow h + X$ can be written as [11, 12]

$$\rho_{\lambda_h \lambda'_h}(h) = \frac{1}{N_h} \sum_{q, X, \lambda_X, \lambda_q, \lambda_{\bar{q}}, \lambda'_q, \lambda'_{\bar{q}}} D_{\lambda_h \lambda_X; \lambda_q, \lambda_{\bar{q}}} \rho_{\lambda_q, \lambda_{\bar{q}}; \lambda'_q, \lambda'_{\bar{q}}}(q\bar{q}) D_{\lambda'_h \lambda_X; \lambda'_q, \lambda'_{\bar{q}}}^*, \quad (11)$$

where $\rho(q\bar{q})$ is the helicity density matrix of the $q\bar{q}$ state created in the annihilation of the unpolarized e^+ and e^- ,

$$\rho_{\lambda_q, \lambda_{\bar{q}}; \lambda'_q, \lambda'_{\bar{q}}}(q\bar{q}) = \frac{1}{4N_{q\bar{q}}} \sum_{\lambda_-, \lambda_+} M_{\lambda_q \lambda_{\bar{q}}; \lambda_- \lambda_+} M_{\lambda'_q \lambda'_{\bar{q}}; \lambda_- \lambda_+}^*. \quad (12)$$

The M 's are the helicity amplitudes for the $e^-e^+ \rightarrow q\bar{q}$ process and the D 's are the fragmentation amplitudes, *i.e.* the helicity amplitudes for the process $q\bar{q} \rightarrow h + X$; the $\sum_{X, \lambda_X}$ stands for the phase space integration and the sum over spins of all the unobserved particles, grouped into a state X . The normalization factors N_h and $N_{q\bar{q}}$ are given by:

$$N_h = \sum_{q, X; \lambda_h, \lambda_X, \lambda_q, \lambda_{\bar{q}}, \lambda'_q, \lambda'_{\bar{q}}} D_{\lambda_h \lambda_X; \lambda_q, \lambda_{\bar{q}}} \rho_{\lambda_q, \lambda_{\bar{q}}; \lambda'_q, \lambda'_{\bar{q}}}(q\bar{q}) D_{\lambda_h \lambda_X; \lambda'_q, \lambda'_{\bar{q}}}^* = \sum_q D_q^h, \quad (13)$$

where D_q^h is the usual fragmentation function of quark q into hadron h , and

$$N_{q\bar{q}} = \frac{1}{4} \sum_{\lambda_q, \lambda_{\bar{q}}; \lambda_-, \lambda_+} |M_{\lambda_q \lambda_{\bar{q}}; \lambda_- \lambda_+}|^2. \quad (14)$$

The helicity density matrix for the $q\bar{q}$ state can be computed in the Standard Model and its non zero elements are given by

$$\rho_{+-;+-}(q\bar{q}) = 1 - \rho_{-+;-+}(q\bar{q}) \simeq \frac{1}{2} \frac{(g_V - g_A)_q^2}{(g_V^2 + g_A^2)_q} \quad (15)$$

$$\rho_{+-;-+}(q\bar{q}) = \rho_{-+;-+}^*(q\bar{q}) \simeq \frac{1}{2} \frac{(g_V^2 - g_A^2)_q}{(g_V^2 + g_A^2)_q} \frac{\sin^2 \theta}{1 + \cos^2 \theta}. \quad (16)$$

These expressions are simple but approximate and hold at the Z_0 pole, neglecting electromagnetic contributions, masses and terms proportional to g_V^l ; the full correct expressions can be found in Ref. [13].

Notice that, inserting the values of the coupling constants

$$g_V^{u,c,t} = \frac{1}{2} - \frac{4}{3} \sin^2 \theta_w \quad g_V^{d,s,b} = -\frac{1}{2} + \frac{2}{3} \sin^2 \theta_w \quad g_A^{u,c,t} = -g_A^{d,s,b} = \frac{1}{2} \quad (17)$$

one has

$$\rho_{+-;-+}(u\bar{u}, c\bar{c}, t\bar{t}) \simeq -0.36 \frac{\sin^2 \theta}{1 + \cos^2 \theta} \quad \rho_{+-;-+}(d\bar{d}, s\bar{s}, b\bar{b}) \simeq -0.17 \frac{\sin^2 \theta}{1 + \cos^2 \theta} . \quad (18)$$

Eq. (18) clearly shows the θ dependence of $\rho_{+-;-+}(q\bar{q})$. In case of pure electromagnetic interactions ($\sqrt{s} \ll M_Z$) one has exactly:

$$\rho_{+-;-+}^\gamma(q\bar{q}) = \frac{1}{2} \frac{\sin^2 \theta}{1 + \cos^2 \theta} . \quad (19)$$

Notice that Eqs. (18) and (19) have the same angular dependence, but a different sign for the coefficient in front, which is negative for the Z contribution.

By using the above equations for $\rho(q\bar{q})$ into Eq. (11) one obtains the most general expression of $\rho(h)$ in terms of the $q\bar{q}$ spin state and the unknown fragmentation amplitudes.

Despite the ignorance of the fragmentation process some predictions can be made [13] by considering the production of hadrons almost collinear with the parent jet: the $q\bar{q} \rightarrow h + X$ fragmentation is then essentially a c.m. forward process and the unknown D amplitudes must satisfy the angular momentum conservation relation [14]

$$D_{\lambda_h \lambda_X; \lambda_q \lambda_{\bar{q}}} \sim \left(\sin \frac{\theta_h}{2} \right)^{|\lambda_h - \lambda_X - \lambda_q + \lambda_{\bar{q}}|} \simeq \left(\frac{p_T}{z\sqrt{s}} \right)^{|\lambda_h - \lambda_X - \lambda_q + \lambda_{\bar{q}}|} , \quad (20)$$

with θ_h the angle between the hadron momentum, $\mathbf{h} = z\mathbf{q} + \mathbf{p}_T$, and the quark momentum $\mathbf{q}$.

The bilinear combinations of fragmentation amplitudes contributing to $\rho(h)$ are then not suppressed by powers of $(p_T/z\sqrt{s})$ only if the exponent in Eq. (20) is zero, which greatly reduces the number of relevant helicity configurations.

The fragmentation process is a parity conserving one and the fragmentation amplitudes must then also satisfy the forward parity relationship

$$D_{-\lambda_h - \lambda_X; -+} = (-1)^{S_h + S_X + \lambda_h - \lambda_X} D_{\lambda_h \lambda_X; +-} . \quad (21)$$

Before presenting analytical and numerical results for the coherent quark fragmentation let us remember that in case of incoherent single quark fragmentation Eq. (11) becomes

$$\rho_{\lambda_h \lambda'_h}(h) = \frac{1}{N_h} \sum_{q, X, \lambda_X, \lambda_q, \lambda'_q} D_{\lambda_h \lambda_X; \lambda_q} \rho_{\lambda_q \lambda'_q} D_{\lambda_h \lambda_X; \lambda_q}^* , \quad (22)$$

where $\rho(q)$ is the quark q helicity density matrix related to $\rho(q\bar{q})$ by

$$\rho_{\lambda_q \lambda'_q} = \sum_{\lambda_{\bar{q}}} \rho_{\lambda_q, \lambda_{\bar{q}}; \lambda'_q, \lambda_{\bar{q}}} (q\bar{q}) . \quad (23)$$

In such a case angular momentum conservation for the collinear quark fragmentation requires $\lambda_q = \lambda_h + \lambda_X$; the Standard Model computation of $\rho(q)$ gives only diagonal terms [$\rho_{++}(q) = \rho_{+-;+-}(q\bar{q})$, $\rho_{--}(q) = \rho_{-+;-+}(q\bar{q})$], and one ends up with the usual probabilistic expression

$$\rho_{\lambda_h \lambda_h}(h) = \frac{1}{N_h} \sum_{q, \lambda_q} \rho_{\lambda_q \lambda_q} D_{q, \lambda_q}^{h, \lambda_h} , \quad (24)$$

where $D_{q, \lambda_q}^{h, \lambda_h}$ is the polarized fragmentation function of a q with helicity λ_q into a hadron h with helicity λ_h . Off-diagonal elements of $\rho(h)$ are all zero.

$e^- e^+ \rightarrow BX$, ($S_B = 1/2$, $p_T/\sqrt{s} \rightarrow 0$)

Let us consider first the case in which h is a spin 1/2 baryon. It was shown in Ref. [12] that in such a case the coherent quark fragmentation only induces small corrections to the usual incoherent description

$$\rho_{++}(B) = \frac{1}{N_B} \sum_q \left[\rho_{+-;+-}(q\bar{q}) D_{q,+}^{B,+} + \rho_{-+;-+}(q\bar{q}) D_{q,-}^{B,+} \right] \quad (25)$$

$$\rho_{+-}(B) = \mathcal{O} \left[\left(\frac{p_T}{z\sqrt{s}} \right) \right] . \quad (26)$$

That is, the diagonal elements of $\rho(B)$ are the same as those given by the usual probabilistic formula (24), with small corrections of the order of $(p_T/z\sqrt{s})^2$, while off-diagonal elements are of the order $(p_T/z\sqrt{s})$ and vanish in the $p_T/\sqrt{s} \rightarrow 0$ limit.

The matrix elements of $\rho(B)$ are related to the longitudinal (P_z) and transverse (P_y) polarization of the baryon:

$$P_z = 2\rho_{++} - 1, \quad P_y = -2\text{Im}\rho_{+-} . \quad (27)$$

Some data are available on Λ polarization, both longitudinal and transverse, from ALEPH Collaboration [15] and they do agree with the above equations. In particular the transverse polarization, at $\sqrt{s} = M_Z$, $p_T \simeq 0.5$ GeV/ c and $z \simeq 0.5$ is indeed of the order 1%, as expected from Eq. (26).

$e^-e^+ \rightarrow V X, (S_V = 1, p_T/\sqrt{s} \rightarrow 0)$

In case of final spin 1 vector mesons one has, always in the limit of small p_T [11], [13]

$$\rho_{00}(V) = \frac{1}{N_V} \sum_q D_{q,+}^{V,0} \quad (28)$$

$$\rho_{11}(V) = \frac{1}{N_V} \sum_q \left[\rho_{+-;+-}(q\bar{q}) D_{q,+}^{V,1} + \rho_{-+;-+}(q\bar{q}) D_{q,-}^{V,1} \right] \quad (29)$$

$$\rho_{1,-1}(V) = \frac{1}{N_V} \sum_{q,X} D_{10;+-} D_{-10;-+}^* \rho_{+-;+-}(q\bar{q}) . \quad (30)$$

Again, the diagonal elements have the usual probabilistic expression; however, there is now an off-diagonal element, $\rho_{1,-1}$, which may survive even in the $p_T/\sqrt{s} \rightarrow 0$ limit. In the sequel we shall concentrate on it. Let us first notice that, in the collinear limit, one has

$$D_{q,+}^{V,0} = \sum_X |D_{0-1;+-}|^2 = D_{q,-}^{V,0} \quad (31)$$

$$D_{q,+}^{V,1} = \sum_X |D_{10;+-}|^2 = D_{q,-}^{V,-1} \quad (32)$$

$$D_{q,-}^{V,1} = \sum_X |D_{12;+-}|^2 = D_{q,+}^{V,-1} , \quad (33)$$

with $D_q^V = D_{q,+}^{V,0} + D_{q,+}^{V,1} + D_{q,+}^{V,-1}$ and $N_V = \sum_q D_q^V$. We also notice that the two fragmentation amplitudes appearing in Eq. (30) are related by parity and their product is always real. ρ_{00} and $\rho_{1,-1}$ can be measured through the angular distribution of two body decays of V .

In order to give numerical estimates of $\rho_{1,-1}$ we make some plausible assumptions

$$D_{q,-}^{h,1} = D_{q,+}^{h,-1} = 0 \quad D_{q,+}^{h,0} = \alpha_q^V D_{q,+}^{h,1} . \quad (34)$$

The first of these assumptions simply means that quarks with helicity $1/2$ ($-1/2$) cannot fragment into vector mesons with helicity -1 ($+1$). This is true for valence quarks assuming vector meson wave functions with no orbital angular momentum, like in $SU(6)$. The second assumption is also true in $SU(6)$ with $\alpha_q^V = 1/2$ for any valence q and V . Rather than taking $\alpha_q^V = 1/2$ we prefer to relate the value of α_q^V to the value of $\rho_{00}(V)$ which can be or has been measured. In fact, always in the $p_T \rightarrow 0$ limit, one has [13]

$$\rho_{00}(V) = \frac{\sum_q \alpha_q^V D_{q,+}^{h,1}}{\sum_q (1 + \alpha_q^V) D_{q,+}^{h,1}} . \quad (35)$$

If α_q^V is the same for all valence quarks in V ($\alpha_q^V = \alpha^V$) one has, for the valence quark contribution:

$$\alpha^V = \frac{\rho_{00}(V)}{1 - \rho_{00}(V)} . \quad (36)$$

Finally, one obtains [13]

$$\rho_{1,-1}(V) \simeq [1 - \rho_{0,0}(V)] \frac{\sum_q D_{q,+}^{V,1} \rho_{+-;-+}(q\bar{q})}{\sum_q D_{q,+}^{V,1}} . \quad (37)$$

We shall now consider some specific cases in which we expect Eq. (37) to hold; let us remind once more that our conclusions apply to spin 1 vector mesons produced in $e^-e^+ \rightarrow q\bar{q} \rightarrow V + X$ processes in the limit of small p_T and large z , *i.e.*, to vector mesons produced in two jet events ($e^-e^+ \rightarrow q\bar{q}$) and collinear with one of them ($p_T = 0$), which is the jet generated by a quark which is a valence quark for the observed vector meson (large z). These conditions should be met more easily in the production of heavy vector mesons.

Among other results one obtains [13]:

$$\rho_{1,-1}(D^*) \simeq [1 - \rho_{0,0}(D^*)] \rho_{+-;-+}(c\bar{c}) \quad (38)$$

$$\rho_{1,-1}(\phi) \simeq [1 - \rho_{0,0}(\phi)] \rho_{+-;-+}(s\bar{s}) \quad (39)$$

$$\rho_{1,-1}(K^{*0}) \simeq \frac{1}{2} [1 - \rho_{0,0}(K^{*0})] [\rho_{+-;-+}(d\bar{d}) + \rho_{+-;-+}(s\bar{s})] . \quad (40)$$

Eqs. (38)-(40) show how the value of $\rho_{1,-1}(V)$ are simply related to the off-diagonal helicity density matrix element $\rho_{+-;-+}(q\bar{q})$ of the $q\bar{q}$ pair created in the elementary $e^-e^+ \rightarrow q\bar{q}$ process; such off-diagonal elements would not appear in the incoherent independent fragmentation of a single quark, yielding $\rho_{1,-1}(V) = 0$.

By inserting into the above equations the value of ρ_{00} when available [16] and the expressions of $\rho_{+-;-+}$, Eq. (18), one has:

$$\rho_{1,-1}(D^*) \simeq -(0.216 \pm 0.007) \frac{\sin^2 \theta}{1 + \cos^2 \theta} \quad (41)$$

$$\rho_{1,-1}(\phi) \simeq -(0.078 \pm 0.014) \frac{\sin^2 \theta}{1 + \cos^2 \theta} \quad (42)$$

$$\rho_{1,-1}(K^{*0}) \simeq -0.170 [1 - \rho_{0,0}(K^{*0})] \frac{\sin^2 \theta}{1 + \cos^2 \theta} . \quad (43)$$

Finally, in case one collects all meson produced at different angles in the full available θ range (say $\alpha < \theta < \pi - \alpha$, $|\cos \theta| < \cos \alpha$) an average should be taken in θ ,

weighting the different values of $\rho_{1,-1}(\theta)$ with the cross-section for the $e^-e^+ \rightarrow V+X$ process; this gives [13]:

$$\langle \rho_{1,-1}(D^*) \rangle_{[\alpha, \pi-\alpha]} \simeq -(0.216 \pm 0.007) \frac{3 - \cos^2 \alpha}{3 + \cos^2 \alpha} \quad (44)$$

$$\langle \rho_{1,-1}(\phi) \rangle_{[\alpha, \pi-\alpha]} \simeq -(0.078 \pm 0.014) \frac{3 - \cos^2 \alpha}{3 + \cos^2 \alpha} \quad (45)$$

$$\langle \rho_{1,-1}(K^{*0}) \rangle_{[\alpha, \pi-\alpha]} \simeq -0.170 [1 - \rho_{0,0}(K^*)] \frac{3 - \cos^2 \alpha}{3 + \cos^2 \alpha} . \quad (46)$$

These results have to be compared with data [16]

$$\rho_{1,-1}(D^*) = -0.039 \pm 0.016 \quad \text{for} \quad z > 0.5 \quad \cos \alpha = 0.9 \quad (47)$$

$$\rho_{1,-1}(\phi) = -0.110 \pm 0.070 \quad \text{for} \quad z > 0.7 \quad \cos \alpha = 0.9 \quad (48)$$

$$\rho_{1,-1}(K^{*0}) = -0.090 \pm 0.030 \quad \text{for} \quad z > 0.3 \quad \cos \alpha = 0.9 \quad (49)$$

which shows a good qualitative agreement with the theoretical predictions. We notice that while the mere fact that $\rho_{1,-1}$ differs from zero is due to a coherent fragmentation of the $q\bar{q}$ pair, the actual numerical values depend on the Standard Model coupling constants; for example, $\rho_{1,-1}$ would be positive at smaller energies, at which the one gamma exchange dominates, while it is negative at LEP energy where the one Z exchange dominates. $\rho_{1,-1}$ has also a peculiar dependence on the meson production angle, being small at small and large angles and maximum at $\theta = \pi/2$. Such angular dependence has been tested in case of K^{*0} production, assuming no dependence of ρ_{00} on θ , and indeed one has [16], in agreement with Eqs. (40) and (18)

$$\left[\frac{\rho_{1,-1}}{1 - \rho_{00}} \right]_{|\cos \theta| < 0.5} \cdot \left[\frac{\rho_{1,-1}}{1 - \rho_{00}} \right]_{|\cos \theta| > 0.5}^{-1} = 1.5 \pm 0.7 \quad (50)$$

These results are encouraging; it would be interesting to have more and more detailed data, possibly with a selection of final hadrons with the required features for our results to hold. It would also be interesting to test the coherent fragmentation of quarks in other processes, like $\gamma\gamma \rightarrow VX$, $pp \rightarrow D^*X$ and $\ell p \rightarrow VX$. The first two processes are similar to $e^-e^+ \rightarrow VX$ in that a $q\bar{q}$ pair is created which then fragments coherently into the observed vector meson; one assumes that the dominating elementary process in $pp \rightarrow D^*X$ is $gg \rightarrow c\bar{c}$. In both these cases one has for $\rho_{+,-;-+}(q\bar{q})$ the same value as in Eq. (19), so that one expects a *positive* value of $\rho_{1,-1}(V)$.

In the last process, the production of vector mesons in DIS, the quark fragmentation is in general a more complicated interaction of the quark with the remnants of the proton and it might be more difficult to obtain numerical predictions. However, if one observes D^* mesons one can assume or select kinematical regions for which the underlying elementary interaction is $\gamma^* g \rightarrow c\bar{c}$: again, one would have the same $\rho_{+-;-+}(c\bar{c})$ as in Eq. (19), and one would expect a positive value of $\rho_{1,-1}(D^*)$. It would indeed be interesting to perform these simple tests of coherent fragmentation effects.

Acknowledgements

I would like to thank the organizers of the Workshop and DESY for financial support

References

- [1] D.L. Adams et al., *Phys. Lett.* **B264** (1991) 462.
- [2] D.L. Adams et al., *Z. Phys.* **C56** (1992) 181.
- [3] J.C. Collins, *Nucl. Phys.* **B396** (1993) 161.
- [4] D. Sivers, *Phys. Rev.* **D41** (1990) 83.
- [5] D. Sivers, *Phys. Rev.* **D43** (1991) 261.
- [6] M. Anselmino, M. Boglione and F. Murgia, *Phys. Lett.* **B362** (1995) 164.
- [7] M. Anselmino, E. Leader and F. Murgia, [hep-ph/9610407](#), to appear in *Phys. Rev.* **D**.
- [8] J.C. Collins, *Nucl. Phys.* **B394** (1993) 169.
- [9] J.C. Collins, S.H. Heppelmann and G.A. Ladinsky, *Nucl. Phys.* **B420** (1994) 565.
- [10] M. Anselmino, A. Drago and F. Murgia, e-Print Archive: [hep-ph/9703303](#).
- [11] M. Anselmino, P. Kroll and B. Pire, *Z. Phys.* **C29** (1985) 135.
- [12] A. Anselm, M. Anselmino, F. Murgia and M. Ryskin, *JETP Lett.* **60** (1994) 496.
- [13] M. Anselmino, M. Bertini, F. Murgia and P. Quintairos, [hep-ph/9704420](#), to appear in *Z. Phys.* **C**.
- [14] See, *e.g.*, C. Bourrely, E. Leader and J. Soffer, *Phys. Rep.* **59** (1980) 95.
- [15] ALEPH Collaboration, *Phys. Lett.* **B374** (1996) 319.
- [16] OPAL Collaboration, [CERN-PPE/97-05](#) and [CERN-PPE/97-094](#).